

DSM 5 Diagnostic Codes Related to Substance Use Disorders

Published on BupPractice (<http://www.buppractice.com>)

For Mental Health Providers- Secondary SUD Dx

[DSM 5 Diagnostic Codes Related to Substance Use Disorders \[1\]](#)

Description: DSM-IV and DSM 5 Diagnostic Codes Related to Substance Use Disorders
(*Note: DSM 5 was released in May 2013 and includes significant changes to diagnosis. For example, it does away with separate "dependence" and "abuse" diagnoses and combines them into "substance use disorder.")

The ICD-10 compliance date is October 1, 2015.

This is a table listing the diagnostic codes for substance use disorders.

| DSM-IV Diagnostic Codes (ICD-9-CM) | DSM-5 Diagnostic Codes (ICD-10-CM) | Substance Use Disorder |
|--|---|------------------------|
| <ul style="list-style-type: none">• 303.00 | <ul style="list-style-type: none">• F10.129• F10.229• F10.929 <p>(Selected code depends on presence and severity of comorbid alcohol use disorder)</p> | Alcohol Intoxication |
| <ul style="list-style-type: none">• 305.00 - Alcohol Abuse• 303.90 - Alcohol Dependence | <ul style="list-style-type: none">• F10.10• F10.20 <p>(Selected code depends on number of symptoms present)</p> | Alcohol Use Disorder |
| <ul style="list-style-type: none">• 304.00 - Opioid Dependence• 305.50 - Opioid Abuse | <ul style="list-style-type: none">• F11.10• F11.20 <p>(Selected code depends on number of symptoms present)</p> | Opioid Use Disorder |
| <ul style="list-style-type: none">• 292.89 | For opioid intoxication without perceptual disturbances: <ul style="list-style-type: none">• F11.129 (with comorbid mild opioid use disorder)• F11.229 (with comorbid moderate or severe opioid use disorder)• F11.929 (with no comorbid opioid use disorder) | Opioid Intoxication |

DSM 5 Diagnostic Codes Related to Substance Use Disorders

Published on BupPractice (<http://www.buppractice.com>)

| | | |
|---|--|--|
| | For opioid intoxication with perceptual disturbances: | |
| | <ul style="list-style-type: none"> • F11.122 (with comorbid mild opioid use disorder) • F11.222 (with comorbid moderate or severe opioid use disorder) • F11.922 (with no comorbid opioid use disorder) | |
| • 292.0 | • F11.23 (Code indicates presence of comorbid moderate or severe opioid use disorder) | Opioid Withdrawal |
| • 292.9 | • F11.99 | Unspecified Opioid-Related Disorder |
| <ul style="list-style-type: none"> • 304.10 - Sedative, Hypnotic, or Anxiolytic Dependence • 305.40 - Sedative, Hypnotic, or Anxiolytic Abuse | <ul style="list-style-type: none"> • F13.10 • F13.20 (Selected code depends on number of symptoms present) | Sedative, Hypnotic, or Anxiolytic Use Disorder |
| <ul style="list-style-type: none"> • 304.20 - Cocaine Dependence • 305.60 - Cocaine Abuse | <ul style="list-style-type: none"> • F14.10 • F14.20 (Selected code depends on number of symptoms present) | Cocaine Use Disorder |
| <ul style="list-style-type: none"> • 304.30 - Cannabis Dependence • 305.20 - Cannabis Abuse | <ul style="list-style-type: none"> • F12.10 • F12.20 (Selected code depends on number of symptoms present) | Cannabis Use Disorder |
| <ul style="list-style-type: none"> • 304.50 - Hallucinogen Dependence • 305.30 - Hallucinogen Abuse | <ul style="list-style-type: none"> • F16.10 • F16.20 (Selected code depends on number of symptoms present) | Other Hallucinogen Use Disorder |
| <ul style="list-style-type: none"> • 305.90 - Inhalent Abuse • 304.60 - Inhalent Dependence | <ul style="list-style-type: none"> • F18.10 • F18.20 (Selected code depends on number of symptoms present) | Inhalent Use Disorder |
| • 304.80 | | Polysubstance |

DSM 5 Diagnostic Codes Related to Substance Use Disorders

Published on BupPractice (<http://www.buppractice.com>)

| | | Dependence |
|---|---|---|
| <ul style="list-style-type: none"> • 304.90 - Other (or Unknown) Substance Dependence • 305.90 - Other (or Unknown) Substance Abuse | <ul style="list-style-type: none"> • F19.10 • F19.20 <p>(Selected code depends on number of symptoms present)</p> | Other (or Unknown) Substance Use Disorder |
| <ul style="list-style-type: none"> • 305.10 - Nicotine Dependence | <ul style="list-style-type: none"> • Z72.0 • F17.200 <p>(Selected code depends on number of symptoms present)</p> | Tobacco Use Disorder |
| <ul style="list-style-type: none"> • 305.70 - Amphetamine Abuse • 304.40 - Amphetamine Dependence | <ul style="list-style-type: none"> • F15.10 • F15.20 <p>(Selected code depends on number of symptoms present)</p> | Amphetamine Use Disorder |
| <ul style="list-style-type: none"> • 305.90 | <ul style="list-style-type: none"> • F15.929 | Caffeine Intoxication |
| <ul style="list-style-type: none"> • 305.90 - Caffeine Intoxication, Inhalent Abuse, or Other (or Unknown) Substance Abuse • 304.60 - Inhalent Dependence | <ul style="list-style-type: none"> • F16.10 • F16.20 <p>(Selected code depends on number of symptoms present)</p> | Phencyclidine Use Disorder |

For a complete list of ICD-10-CM codes, visit the World Health Organization's website.

Links:

[1] <http://www.buppractice.com/node/2633>